



1. (a) Let a directed graph G_1 be given.



Does each of the following list of vertices form a path in G_1 ? If yes, determine (by circling) if the path is simple, if it is a circuit, and give its length.

- | | |
|------------------|---|
| a, b, e, c, b | Yes [simple circuit length <input type="text"/>] No |
| a, d, a, d, a | Yes [simple circuit length <input type="text"/>] No |
| a, d, e, b, a | Yes [simple circuit length <input type="text"/>] No |
| a, b, e, c, b, a | Yes [simple circuit length <input type="text"/>] No |

(b) For the simple graph G_2




Find M^2 , where M is the adjacency matrix of G_2

$$M^2 = \left\{ \begin{array}{cccc} \square & \square & \square & \square \\ \square & \square & \square & \square \\ \square & \square & \square & \square \\ \square & \square & \square & \square \end{array} \right\}$$


Find the number of paths from A to D in G_2 of length 2.

2. Let $f(n) = 1 + 2 + 3 + \dots + n$. Show that $f(n)$ is $O(n^2)$. Be sure to specify the values of the witnesses C and k .

3. Determine whether the given pair of graphs is isomorphic. Exhibit an isomorphism or provide a rigorous argument that none exists.



G




H

4. Show that a vertex c in the connected simple graph G is a cut vertex if and only if there are vertices u and v , both different from c , such that every path between u and v passes through c .

5.


(a) Is there an Euler circuit in the following graph? If so, find such a circuit. If not, explain why no such circuit exists.

(b) Is there a Hamilton circuit in the following graph? If so, find such a circuit. If not, prove why no such circuit exists.



6. Let $f(n) = 2n^2 + 5n\log(n) + 8n + 7$. Show that $f(n)$ is $O(n^2)$. Be sure to specify the values of the witnesses C and k .

7. Use Dijkstra's algorithm to find the length of the shortest path between the vertices a and z in the following weighted graph.



a	b	c	d	e	z	S
0	∞	∞	∞	∞	∞	a
X						
X						
X						
X						
X						
X						
X						
X						

8. How many vertices and how many edges does each of the following graphs have?

(a) K_5

(b) C_4

(c) W_5

(d) $K_{2,5}$

9. Describe an algorithm for finding the second largest integer in a sequence of distinct integers. Give a big-O estimate of the number of comparisons used by your algorithm.

10. Give a recursive algorithm for finding the string w^i , the concatenation of i copies of w , when w is a bit string.

- 11.** Give a recursive definition of
- a) the set of odd positive integers.
 - b) the set of positive integer powers of 3.
 - c) the set of polynomials with integer coefficients.

- 12.** Let S be the subset of the set of ordered pairs of integers defined recursively by
Basis step: $(0, 0) \in S$.
Recursive step: If $(a, b) \in S$, then $(a + 2, b + 3) \in S$ and $(a + 3, b + 2) \in S$.
- a)** List the elements of S produced by the first two applications of the recursive definition.
 - b)** Use structural induction to show that $5 \mid a + b$ when $(a, b) \in S$.

13. Write a pseudocode for an algorithm for evaluating a polynomial of degree n ,

$$p(x) = a_n x^n + a_{n-1} x^{n-1} + \dots + a_1 x + a_0, \text{ at } x = c.$$

What is big-O estimate of the time complexity of your algorithm (in terms of the number of multiplications and additions used) as a function of n ? Explain your answer.

14. Show that $\log(n!)$ is $\Theta(n \cdot \log(n))$.

15. Prove that $1 \cdot 1! + 2 \cdot 2! + \dots + n \cdot n! = (n + 1)! - 1$ whenever n is a positive integer.

16. Prove that $1 + 3 + \dots + (2n - 1) = n^2$ whenever n is a positive integer.