



1. (a) Let a directed graph G_1 be given.



Does each of the following list of vertices form a path in G_1 ? If yes, determine (by circling) if the path is simple, if it is a circuit, and give its length.

- | | |
|------------------|---|
| a, b, e, c, b | Yes [simple circuit length <input type="text"/>] No |
| a, d, a, d, a | Yes [simple circuit length <input type="text"/>] No |
| a, d, e, b, a | Yes [simple circuit length <input type="text"/>] No |
| a, b, e, c, b, a | Yes [simple circuit length <input type="text"/>] No |

(b) For the simple graph G_2





Find M^2 , where M is the adjacency matrix of G_2

$$M^2 = \left\{ \begin{array}{cccc} \square & \square & \square & \square \\ \square & \square & \square & \square \\ \square & \square & \square & \square \\ \square & \square & \square & \square \end{array} \right\}$$

Find the number of paths from A to D in G_2 of length 2.

2. Provide a pseudo code of an algorithm for finding a closest pair of numbers in a set of n real distinct numbers and give a worst-case estimate of the number of comparisons.

3. Determine whether the given pair of graphs is isomorphic. Exhibit an isomorphism or provide a rigorous argument that none exists.




4. Let $a_1 = 2$, $a_2 = 9$, and $a_n = 2a_{n-1} + 3a_{n-2}$ for $n \geq 3$. Show using induction that $a_n \leq 3^n$ for all positive integers n .

5. Use mathematical induction to show that $\sum_{j=0}^n (j + 1) = \frac{(n+1)(n+2)}{2}$ whenever n is a nonnegative integer.


6. Let $f(n) = 2n \log(n^2 + 5) + 3n + 1$. What is big-O estimate of $f(n)$? Be sure to specify the values of the witnesses C and k .

7. Use Dijkstra's algorithm to find the length of the shortest path between the vertices a and z in the following weighted graph. Use the table below to log in your computation.



a	b	c	d	e	z	S
0	∞	∞	∞	∞	∞	a
X						
X						
X						
X						
X						
X						
X						
X						
X						

Draw a tree representing the shortest distances from a to each of the other vertices. Indicate the distance next to each vertex.



8. How many vertices and how many edges does each of the following graphs have?

(a) K_5

(b) C_4

(c) W_5

(d) $K_{2,5}$

9. Write a pseudocode for an algorithm for evaluating a polynomial of degree n ,

$$p(x) = a_n x^n + a_{n-1} x^{n-1} + \dots + a_1 x + a_0, \text{ at } x = c.$$

What is big-O estimate of the time complexity of your algorithm (in terms of the number of multiplications and additions used) as a function of n ? Explain your answer.

10. Let S be the subset of the set of ordered pairs of integers defined recursively by

Basis step: $(0, 0) \in S$.

Recursive step: If $(a, b) \in S$, then $(a + 2, b + 3) \in S$ and $(a + 3, b + 2) \in S$.

a) List the elements of S produced by the first two applications of the recursive definition.

b) Use structural induction to show that $5 \mid a + b$ when $(a, b) \in S$.

11. For which values of n do these graphs have an Euler circuit?
a) K_n b) C_n c) W_n d) Q_n

12. What is the effect in the time required to solve a problem when you double the size of the input from n to $2n$? Express your answer in the simplest form possible, either as a ratio or a difference. Explain the meaning of your answer.
a) $\log n$
b) $100n$
c) n^2

13. Use mathematical induction to prove that every postage of n cents greater than 5 cents can be formed from 3-cent and 4-cent stamps.

14. Give a recursive algorithm for finding the maximum of a finite set of integers, the recursion should make use of the fact that the maximum of n integers is the larger of the last integer in the list and the maximum of the first $n - 1$ integers in the list.